

Subfactor projections

Mladen Bestvina and Mark Feighn*

January 26, 2013

Abstract

When two free factors A, B of a free group $\mathbb{F}_n$ are in “general position” we define the projection $\pi_A(B)$ of B to the splitting complex (alternatively, the complex of free factors) $\mathcal{S}(A)$ of A . We show that the projections satisfy properties analogous to subsurface projections introduced by Masur and Minsky. We use the subfactor projections to construct an action of $\text{Out}(\mathbb{F}_n)$ on a finite product of hyperbolic spaces where every automorphism with exponential growth acts with positive translation length. We also prove a version of the Bounded geodesic image theorem. In the appendix, we give a sketch of the proof of the Handel-Mosher hyperbolicity theorem for the splitting complex using (liberal) folding paths.

1 Introduction

The work of Howard Masur and Yair Minsky [12, 13] has served as an inspiration in the study of $\text{Out}(\mathbb{F}_n)$, the outer automorphism group of the free group $\mathbb{F}_n$ of rank n . Their work starts by proving that the curve complex associated to a compact surface is hyperbolic. There are several established analogs of the curve complex in the $\text{Out}(\mathbb{F}_n)$ setting that have recently been shown to be hyperbolic, see [3, 7, 11] and [8, Section 8].

The complex $\mathcal{F}$ of free factors is the simplicial complex whose vertices are conjugacy classes of nontrivial free factors of $\mathbb{F}_n$ and a simplex is formed by those vertices that have representatives that can be arranged in a chain. (A subgroup A of $\mathbb{F}_n$ is a nontrivial free factor if $\mathbb{F}_n = A * B$ for another subgroup B with $A \neq 1 \neq B$.)

The complex $\mathcal{S}$ of free splittings, or the *sphere complex*, or the simplicial completion of Outer space, is the simplicial complex whose k -simplices are

*Both authors gratefully acknowledge the support by the National Science Foundation.

$(k + 1)$ -edge free splittings (nontrivial minimal reduced simplicial $\mathbb{F}_n$ -trees with trivial edge stabilizers and $(k + 1)$ orbits of edges), up to equivariant isomorphism.

Theorem 1.1 ([3]). *$\mathcal{F}$ is hyperbolic.*

Theorem 1.2 (Handel-Mosher [7]). *$\mathcal{S}$ is hyperbolic.*

Kapovich-Rafi [9] show how to derive Theorem 1.1 from Theorem 1.2 in a simple way, and Hilion-Horbez [8] give a simpler proof of Theorem 1.2 using surgery paths.

Masur and Minsky also define *subsurface projections*. If α is a simple closed curve in a surface S and A is a subsurface such that α and A cannot be made disjoint by an isotopy (we also say they *intersect*), there is a curve $\alpha|A$ in the curve complex of A , well defined up to a bounded set, obtained by “closing up the intersection of α and A along the boundary of A ”. If A, B are two subsurfaces that intersect, one defines the projection $\pi_A(B)$ of B to A by taking $\partial B|A$. We view $\pi_A(B)$ as a bounded subset of the curve complex of A . Masur and Minsky use subsurface projections to estimate distances in mapping class groups, and more generally, to understand their large-scale geometry. To encapsulate this, we first recall a theorem.

Theorem 1.3 ([2]). *Let $\mathcal{Y}$ be a collection of δ -hyperbolic spaces and for every pair $A, B \in \mathcal{Y}$ of distinct elements suppose we are given a uniformly bounded subset $\pi_A(B) \subset A$, called the projection of B to A . Denoting by $d_A(B, C)$ the diameter of $\pi_A(B) \cup \pi_A(C)$, assume the following holds: there is a constant $K > 0$ such that*

- *if $A, B, C \in \mathcal{Y}$ are distinct, at most one of 3 numbers*

$$d_A(B, C), d_B(A, C), d_C(A, B)$$

is $> K$, and

- *for any distinct A, B the set*

$$\{C \in \mathcal{Y} - \{A, B\} \mid d_C(A, B) > K\}$$

is finite.

Then there is a hyperbolic space Y and an isometric embedding of each $A \in \mathcal{Y}$ onto a convex set in Y so the images are pairwise disjoint and the nearest point projection of any B to any $A \neq B$ is within uniformly bounded distance of $\pi_A(B)$. Moreover, the construction is equivariant with respect to any group acting on $\mathcal{Y}$ by isometries.

In the setting of subsurface projections, $\mathcal{Y}$ will be the collection of curve complexes associated to a collection of pairwise intersecting subsurfaces. That both bullets hold in the setting of subsurface projections (whenever they are both defined) is the work of Masur-Minsky [12, 13] and Behrstock [1]. There is a simple proof of the first bullet due to Leininger, see [10], and of the second bullet in [2].

It is further shown in [2] that there is a finite equivariant coloring of the collection of all subsurfaces of S , and the boundaries of two distinct subsurfaces with the same color intersect. Applying Theorem 1.3 to each color separately and then taking the cartesian product one obtains:

Theorem 1.4 ([2]). *The mapping class group of S acts by isometries on the finite product $Y_1 \times \cdots \times Y_k$ of hyperbolic spaces. Moreover, orbit maps are quasi-isometric embeddings.*

The second statement in the theorem recasts the Masur-Minsky distance estimate.

The goal of this paper is to take a step in this direction in the $Out(\mathbb{F}_n)$ setting. If A and B are distinct conjugacy classes of nontrivial free factors, we attempt to define $\pi_A(B)$, a bounded subset of the splitting complex $\mathcal{S}(A)$ of A . We start with a simplicial $\mathbb{F}_n$ -tree T representing a splitting of $\mathbb{F}_n$ where B fixes a vertex. Restricting the action to A and passing to the minimal subtree (assuming A does not fix a vertex of T) gives a splitting of A . Different choices of T will in general produce unbounded sets in $\mathcal{S}(A)$, but when A and B are in “general position” the ambiguity is bounded.

Main Theorem. *Assume $n = \text{rank}(\mathbb{F}_n) \geq 3$. The set of vertices of $\mathcal{F}$ can be equivariantly colored in finitely many colors so that when A and B are distinct and either have the same color or are at distance > 4 then the projection $\pi_A(B)$ is well defined as a bounded subset of $\mathcal{S}(A)$. Moreover, restricting to one color, both bullets in Theorem 1.3 hold.*

The Main Theorem is proved in Section 4. We thus obtain an isometric action of $Out(\mathbb{F}_n)$ on a finite product of hyperbolic spaces. However, orbit maps are not quasi-isometric embedding. We discuss this further in Section 6.

Theorem 5.1. *$Out(\mathbb{F}_n)$ acts on a finite product $Y = Y_1 \times \cdots \times Y_k$ of hyperbolic spaces so that every exponentially growing automorphism has positive translation length.*

The key to understanding the projections is the analysis of a path in

Outer space $\mathcal{X}(A)$ of A induced by a path G_t in $\mathcal{X}$. Here is a motivating example. For a review of the terminology, see Section 2.

Consider the automorphism Φ of $\mathbb{F}_3 = \langle x, y, z \rangle$ given by $\phi : \mathbb{F}_3 \rightarrow \mathbb{F}_3, x \mapsto y, y \mapsto z$ and $z \mapsto zx$. The map on the rose R_0 with loops representing x, y, z is a train track map for Φ . The vertex has 4 gates, namely $\{x, y, z\}, \{X\}, \{Y\}$ and $\{Z\}$ (the direction oriented the same way as the edge x is also called x ; the direction pointing the opposite way is X etc). The dilatation λ of Φ satisfies $\lambda^3 - \lambda^2 - 1 = 0$ and $\lambda = 1.46557 \dots$. The lengths of edges x, y, z are proportional to $1, \lambda, \lambda^2$.

Let $A = \langle x, y \rangle$. We are interested in picturing the core of the covering space corresponding to A of the graphs $R_k = R_0 \Phi^k$, which we view as a discrete path in $\mathcal{X}$ as $k \in \mathbb{Z}$ (but one can extend it to an axis $R_k, k \in \mathbb{R}$ of Φ). This core has an induced metric and the structure of legal and illegal turns. Since $\ell_{R_0 \Phi^k}(\gamma) = \ell_{R_0}(\Phi^k(\gamma))$, this amounts to picturing the $\phi^k(A)$ -cover of R_0 . See Figure 1.




Figure 1: $A|R_0\Phi^k$ for $-5 \leq k \leq 7$.

For example, for $k = 2$ we compute $\phi^2(A) = \langle z, xz \rangle = \langle x, z \rangle$, and for $k = 6$ we get $\phi^6(A) = \langle zxyzzx, zxyzzxzy \rangle = \langle zxy, zzx \rangle$ which is represented by a theta graph with the middle edge z going up, the left edge xy and the right edge zx , both going down. The illegal turn is at the top.

The reader should note the following features:

- As $k \rightarrow \infty$ the volume of $A|R_k$ goes to ∞ and in fact the topological edges of $A|R_k$ are legal and grow exponentially.
- As $k \rightarrow -\infty$ the volume of $A|R_k$ goes to ∞ . The topological edges consist of an exponentially growing number of legal segments of length < 2 and the turns between consecutive such segments are illegal.

Our two main technical lemmas restrict the interval where $A|R_k$ can make progress in $\mathcal{S}(A)$. Lemma 3.1 says:

- If $A|R_k$ has an illegal turn in a topological edge or a vertex with one gate, then the image of $A|R_t$, $t \in (-\infty, k]$, in $\mathcal{S}(A)$ is uniformly bounded.
- If $A|R_k$ has a legal segment of length 2 in a topological edge, then the image of $A|R_t$, $t \in [k, \infty)$, in $\mathcal{S}(A)$ is uniformly bounded.

An easy computation shows that the first bullet holds for $k = -1$ and the second for $k = 9$. Thus all the progress in $\mathcal{S}(A)$ is restricted to the interval $[-1, 9]$. Lemma 4.4 shrinks this interval further: if every edge of R_k has at least two lifts to $A|R_k$ then the image of $A|R_t$, $t \in [k, \infty)$, in $\mathcal{S}(A)$ is uniformly bounded. In our example this holds for $k = 7$.

To simplify the analysis, we will work with *greedy* folding paths as we did in [3]. The above path (an axis of Φ) is *not* a greedy folding path since not all illegal turns are folding. For example, we will argue in the proof of Lemma 3.1 that if $A|G_{t_0}$ has an interior illegal turn then the same is true for all $A|G_t$ for $t < t_0$. Note that this fails for our path R_k .

We conclude the paper with the Bounded geodesic image theorem, analogous to the theorem of Masur and Minsky [13].

Theorem 5.3 (Bounded geodesic image theorem). *Let (A_i) be a geodesic in $\mathcal{F}$ and $A \in \mathcal{F}$ at distance > 4 from each A_i . Then the set of projections of the A_i 's to $\mathcal{S}(A)$ is a uniformly bounded set.*

Sabalka and Savchuk in [14] define a different notion of projection in the language of sphere systems. It is a partially defined map from the sphere complex of $M = \#_1^n S^1 \times S^2$ to the sphere complex of a submanifold of M which is a complementary component of a sphere system. They also prove versions of the Behrstock inequality and the Bounded geodesic image theorem. It would be interesting to understand the precise relationship between the two notions of projection.

A technical tool we use is that folding paths in Outer space project to reparametrized quasi-geodesics in the splitting complex $\mathcal{S}$. In [7] Handel and Mosher use a special kind of folding paths (maximal folds on maps with at least 3 gates at every vertex) that does not suffice for our purposes. In the appendix we follow the same outline as in [7] but we use (liberal) folding paths. Interestingly, the proof becomes simpler since we do not have to arrange the 3 gates condition after every modification.

Acknowledgements. We thank Ken Bromberg, Ursula Hamenstädt, Ilya Kapovich and Patrick Reynolds for their interest and useful comments.

2 Review

In this section we recall basic definitions and facts about Outer space with an emphasis on folding paths. For more information see [3].

A *graph* is a 1-dimensional CW complex. The *rose* in rank n is the graph R_n with 1 vertex and n edges. A finite connected graph is a *core graph* if it has no valence 1 vertices. The *core* of a connected graph Γ with finitely generated $\pi_1(\Gamma)$ is the unique core graph $C \subset \Gamma$ so that inclusion is a homotopy equivalence.

A *marking* of a graph Γ is a homotopy equivalence $f : R_n \rightarrow \Gamma$. A *metric* on a graph Γ is a function that assigns a positive number $\ell(e)$, called *length*, to each edge e of Γ . A metric allows one to view Γ as a path metric space. Culler-Vogtmann's *Outer space* $\mathcal{X}$ is the space of equivalence classes of triples (Γ, ℓ, f) where Γ is a finite graph with no vertices of valence 1 or 2, f is a marking, and ℓ is a metric on Γ with volume (i.e. sum of the lengths of all edges) equal to 1. Two such triples (Γ, ℓ, f) and (Γ', ℓ', f') are equivalent if there is an isometry $\phi : \Gamma \rightarrow \Gamma'$ such that $\phi f \simeq f'$ (homotopic).

The group $Out(\mathbb{F}_n)$ can be identified with the group of homotopy equivalences on R_n . The action of $Out(\mathbb{F}_n)$ on $\mathcal{X}$ (on the right) is defined by changing the marking: $[(\Gamma, \ell, f)] \cdot \Phi = [(\Gamma, \ell, f\Phi)]$.

We frequently simplify the notation and replace the equivalence class $[(\Gamma, \ell, f)]$ by Γ .

The set $\mathcal{C}$ of nontrivial conjugacy classes in $\mathbb{F}_n$ can be identified with the set of homotopy classes of immersed loops in R_n . If $\alpha \in \mathcal{C}$ and $[\Gamma, \ell, f] \in \mathcal{X}$, we define $\alpha|\Gamma$ to be the immersed loop homotopic to $f(\alpha)$, and we let $\ell_\Gamma(\alpha)$ be the length of $\alpha|\Gamma$. We then have $\ell_{\Gamma\Phi}(\alpha) = \ell_\Gamma(\Phi(\alpha))$.

If $\Gamma, \Gamma' \in \mathcal{X}$ we say that a map $\phi : \Gamma \rightarrow \Gamma'$ is a *difference of markings* if $\phi f \simeq f'$. We will normally consider only maps that are linear (constant speed) on each edge. Denote by $\sigma(\phi)$ the Lipschitz constant of ϕ , i.e. the

maximum speed over all edges of Γ . The Lipschitz distance is defined by

$$d(\Gamma, \Gamma') = \min_{\phi} \log \sigma(\phi)$$

where the minimum is taken over all difference of markings maps $\phi : \Gamma \rightarrow \Gamma'$.

A *train track structure* on a graph Γ consists of an equivalence relation on the set of directions out of each vertex of Γ . One usually pictures graphs with a train track structure so that equivalent directions are tangent. Equivalence classes are called *gates*. The tension graph Δ_{ϕ} has a natural train track structure, where two directions d_1, d_2 out of a vertex $v \in \Delta_{\phi}$ are equivalent if they are mapped by ϕ to the same direction out of $\phi(v)$ (which is not required to be a vertex). We say ϕ is *optimal* if $\Delta_{\phi} = \Gamma$ and there are at least two gates at every vertex of Γ . A turn (i.e. a pair of distinct directions out of a vertex) is *illegal* if the directions are in the same gate, and otherwise it is *legal*. An immersed path or a loop is *legal* if it takes only legal turns.

An optimal map $\phi : \Gamma \rightarrow \Gamma'$ is a *train track map* if there is a train track structure on Γ which is preserved by ϕ , in the sense that legal paths are mapped to legal paths.

Now let $\phi : \Gamma \rightarrow \Gamma'$ be optimal. We define a (*greedy*) *folding path* induced by ϕ as follows. It is the unique (up to reparametrization) path G_t , $t \in [\alpha, \omega]$ with $G_{\alpha} = \Gamma$, $G_{\omega} = \Gamma'$, and for every $t \in [\alpha, \omega)$ there is $\epsilon > 0$ such that for every $\tau \in [t, t + \epsilon]$ the graph G_{τ} is obtained from G_t by identifying segments of length $\tau - t$ leaving any vertex in equivalent directions and then rescaling to restore volume 1. We normally parametrize the path by arclength, i.e. so that for $t_1 < t_2$ we have $d(G_{t_1}, G_{t_2}) = t_2 - t_1$. There are induced optimal maps $G_{t_1} \rightarrow G_{t_2}$ for $t_1 < t_2$ and they compose naturally.

If $\phi : G \rightarrow G'$ is any difference of markings map there is an *induced* path from G to G' as follows. It is a concatenation of a linear path within a simplex from $G \rightarrow G''$ and a folding path from G'' to G' induced by an optimal map (see [6], also [3, Proposition 2.7]).

Let G_t be a folding path defined on a closed interval $[\alpha, \omega]$. Then the interval can be subdivided into finitely many subintervals $[t_i, t_{i+1}]$ so that for every $t \in (t_i, t_{i+1})$ all illegal turns are at vertices with two gates, with one gate consisting of a single direction. Moreover, all G_t in this interval are canonically homeomorphic with the homeomorphism preserving the train track structure. As t increases in this open interval, G_t moves along a straight line in the interior of a simplex in $\mathcal{X}$, i.e. only the lengths of the edges change. See Figure 2.

The proof of these facts follows from the analysis of folding paths in [3]. We understand what happens to the right of a given G_t , by the definition




Figure 2: Folding path between events

of a folding path. To understand what happens on an interval $[t_0 - \epsilon, t_0]$ we use the local description of unfolding in [3, Section 4].

Let N be a small connected neighborhood of a vertex v in G_{t_0} . Thus N is a wedge of arcs along their endpoints, with v the wedge point. For small $\epsilon > 0$ the preimage $N_\epsilon \subset G_{t_0 - \epsilon}$ of N is a tree that comes with a height function and a collection of *widgets*. See Figure 3.




Figure 3: The structure of N_ϵ . This one has 3 widgets and 5 vertices at height 0.

The preimages of v are at height 0. All illegal turns are at the vertices at height ϵ . Each of these vertices has two gates, one gate consists of a single direction drawn upwards and the other gate has ≥ 2 directions all of which fold together at $t = t_0$. The union of the edges in the downwards gate at each vertex at height ϵ is a widget. The union of all widgets is a tree.

We also recall that there are coarse projection maps $\mathcal{X} \rightarrow \mathcal{S} \rightarrow \mathcal{F}$ defined by sending a graph $\Gamma \in \mathcal{X}$ to a 1-edge splitting of $\mathbb{F}_n$ obtained by collapsing all but one edge of Γ , and sending a 1-edge splitting to a vertex stabilizer. A useful fact, supplementing Theorems 1.1 and 1.2, is that the image of a folding path in $\mathcal{X}$ is a reparametrized quasi-geodesic in both $\mathcal{F}$ and $\mathcal{S}$, with uniform constants (see [3] for $\mathcal{F}$ and the appendix of this paper for $\mathcal{S}$).

3 Folding path lemmas

Recall that $A|G$ denotes the core of the covering space of G corresponding to a subgroup A of $\mathbb{F}_n$. In this section we prove two technical lemmas about induced paths. The setting in both is that G_t is a folding path in $\mathcal{X}$, $A|G_t$ is the induced path in Outer space $\mathcal{X}(A)$ of A , and we are interested in identifying geometrically the portions of the path $A|G_t$ where it (possibly) makes progress in the splitting complex $\mathcal{S}(A)$ of A .

Convention. For the rest of this paper we use *bounded* to mean *bounded by a function of $\text{rank}(\mathbb{F}_n)$ alone*.

By a *topological* (or *natural*) *edge* in a core graph we mean an edge with respect to the cell structure with all vertices of valence ≥ 3 , and likewise, vertices with valence ≥ 3 are *natural* (when the core graph is a circle we don't talk about topological edges). An *interior illegal turn* is the turn (pair of directions) at a valence two vertex with one gate. Such turns arise in $A|G_t$.

Lemma 3.1. *Given a folding path G_t , $t \in [\alpha, \omega]$, and a f.g. subgroup $A < \mathbb{F}_n$, the path $A|G_t$ can be subdivided into 3 segments $[\alpha, \beta)$, $[\beta, \gamma)$, $[\gamma, \omega]$ (with one or two segments possibly omitted) so that:*

- (1) *On the first segment $[\alpha, \beta)$ $A|G_t$ has a natural vertex with one gate or an interior illegal turn.*
- (2) *On the middle segment $[\beta, \gamma)$ $A|G_t$ has all edges of size ≤ 2 , hence it has bounded volume, all vertices have ≥ 2 gates and there are no interior illegal turns.*
- (3) *On the last segment $[\gamma, \omega]$ $A|G_t$ has a legal segment of length ≥ 2 contained in a topological edge.*

Moreover, the image of the first and the last segment in the splitting complex $\mathcal{S}(A)$ of A is bounded.

Proof. Denote by J the set of those $t_0 \in [\alpha, \omega]$ such that $A|G_{t_0}$ contains a vertex with one gate or an interior illegal turn. We will first argue that J is open in $[\alpha, \omega]$. It is clear from the definition of folding paths that if $t_0 \in J$ and $t_0 \neq \omega$ then $[t_0, t_0 + \epsilon) \subset J$ for small ϵ . Indeed, the effect of local folding at a valence k vertex of $A|G_{t_0}$ with one gate is that an edge with a vertex of valence 1 develops, and after removing it (which is required when passing to the core) a valence k vertex with one gate remains. Assume now that

$t_0 \in J$ and $t_0 \neq \alpha$. We will show that for sufficiently small $\epsilon > 0$ we have $[t_0 - \epsilon, t_0] \subset J$ and this will complete the proof that J is open. The argument uses the description of unfolding in Section 2 and will at the same time lay the groundwork for showing that on J the path is not making progress in $\mathcal{S}(A)$.

Let v be a vertex of $A|G_{t_0}$ of valence k and with one gate (so $k = 2$ in the case of an interior illegal turn). There are k free splittings of $A|G_{t_0}$ obtained by collapsing all edges of $A|G_{t_0}$ except for one edge incident to v (some of these splittings may coincide). These splittings are always uniformly close in $\mathcal{S}(A)$. We will further argue that either

- (A1) $A|G_{t_0-\epsilon}$ also has a vertex of valence k and one gate and the associated splittings are the same as at time t_0 , or
- (A2) $A|G_{t_0-\epsilon}$ has a vertex of valence $< k$ and one gate and the associated splittings are uniformly close to the ones at time t_0 (this is of course possible only if $k > 2$).

Let N , N_ϵ be as in Section 2, i.e. N is a small neighborhood of the image of v in G_{t_0} and N_ϵ is the preimage of N in $G_{t_0-\epsilon}$. Denote by $\tilde{G}_t$ the covering of G_t corresponding to A ; thus $A|G_t \subset \tilde{G}_t$ is the core. Let $\tilde{N}$ be the lift of N to $\tilde{G}_{t_0}$ that contains v and $\tilde{N}_\epsilon$ the preimage of $\tilde{N}$ under the lifted folding map $\tilde{G}_{t_0-\epsilon} \rightarrow \tilde{G}_{t_0}$. Since $\tilde{N}_\epsilon \rightarrow \tilde{N}$ is a homeomorphism, we will talk about height, widgets, etc in $\tilde{N}_\epsilon$. All edges in a widget in $\tilde{N}_\epsilon$ map to the same small segment in $\tilde{G}_{t_0}$ with one endpoint v . If this segment is in the core $A|G_{t_0}$ we will call the widget *essential*, and otherwise it is *inessential*. Likewise, an edge below height 0 is essential if its image is in $A|G_{t_0}$, otherwise it is inessential. Altogether, there are precisely k essential edges and widgets. We now observe:

- (B1) If x is a vertex of $\tilde{N}_\epsilon$ at height ϵ that belongs to an inessential widget, then the edge that goes upwards from x does not belong to $A|G_{t_0-\epsilon}$.
- (B2) Essential widgets and essential edges are pairwise disjoint.
- (B3) If z is a vertex of $\tilde{N}_\epsilon$ at height 0 that belongs to an essential widget, then any edges that go down from z do not belong to $A|G_{t_0-\epsilon}$.

All of these statements are consequences of the facts that v has only one gate in $A|G_{t_0}$ and that the turns at z and at x involving the upward edge are legal.

Essential edges below height 0 are the only ones that map to their respective segments adjacent to v so they are necessarily part of the core $A|G_{t_0-\epsilon}$.

For the same reason, at least one edge in each essential widget belongs to $A|G_{t_0-\epsilon}$. Denote by $\tilde{N}'_\epsilon$ the hull of all essential edges and widgets, together with edges above essential widgets. The rest of $\tilde{N}_\epsilon$ is not part of the core (and even some of the edges in essential widgets might be outside the core). We retain the terminology from $\tilde{N}_\epsilon$, e.g. an inessential widget in $\tilde{N}'_\epsilon$ is the intersection of an inessential widget in $\tilde{N}_\epsilon$ with $\tilde{N}'_\epsilon$. But note that in $\tilde{N}'_\epsilon$:

- (C1) There are no edges above inessential widgets.
- (C2) There are no edges below essential widgets.
- (C3) Each inessential widget contains at least 2 edges.
- (C4) $\tilde{N}'_\epsilon$ must contain at least one inessential widget.
- (C5) Each complementary component of the height ϵ vertex in each inessential widget contains at least one essential edge or essential widget.
- (C6) Every inessential widget has $\leq k$ edges.

(C4) follows from the fact that essential widgets are pairwise disjoint and it's impossible for $\tilde{N}'_\epsilon$ to consist just of a single essential widget. (C5) follows from the construction of $\tilde{N}'_\epsilon$, and (C6) follows from (C5).

If $\tilde{N}'_\epsilon$ has an inessential widget with $< k$ edges, we are done (the associated splittings are clearly within bounded distance of the original k splittings). Suppose W is an inessential widget in $\tilde{N}'_\epsilon$ with exactly k edges. We will argue that the k splittings associated to this widget are exactly the same as the original k splittings. All other inessential widgets in $\tilde{N}'_\epsilon$ have two edges (see (C5)). Thus each edge of W is attached to a (possibly empty) sequence of inessential widgets with 2 edges and then either an essential edge or an essential widget. The map $\tilde{N}'_\epsilon \rightarrow \tilde{N}$ factors through $\tilde{N}''_\epsilon$ obtained from $\tilde{N}'_\epsilon$ by folding together all edges in each essential widget. The folding map does not affect the associated splittings of A . Now observe that $\tilde{N}''_\epsilon$ is a k -pronged star (wedge of k arcs along their endpoints) and the map $\tilde{N}''_\epsilon \rightarrow \tilde{N}'_\epsilon$ amounts to partially folding together the k prongs and this clearly does not affect the splittings.

In Figures 4, 5 and 6 we illustrate the 3 possibilities when $k = 2$.

We now know that J is open. Next, we observe that J must be an initial segment of $[\alpha, \omega]$ (or empty). Indeed, otherwise we would have $t_0 \notin J$, but $(t_0, t_0 + \epsilon) \subset J$. However, a local fold at $A|G_{t_0}$ cannot produce a vertex with one gate. Similarly, splittings visible at $t_0 \in J$ are also visible on $[t_0, t_0 + \epsilon]$. The picture is that as $A|G_t$ unfolds (with t decreasing), our valence k vertex




Figure 4: This is $\tilde{N}'_\epsilon$ for the example in Figure 3 assuming there are two essential edges, the leftmost and the rightmost.




Figure 5: This is N'_ϵ for the example in Figure 3 assuming that the leftmost edge and the rightmost widget are essential.

with one gate undergoes (perhaps simultaneously) a sequence of changes where it lowers the valence, but retains the feature that it has only one gate. Between these events, the associated splittings do not change, and at the events they change but only by a bounded distance.

Now (A1), (A2) and this discussion then prove that we can take $[\alpha, \beta) = J$ and the image of G_t , $t \in J$ is bounded in $\mathcal{S}(A)$.

To conclude the proof of Lemma 3.1 we have to consider the last segment. Suppose $A|G_{t_0}$ has a legal segment of length ≥ 2 in a topological edge. Consider the splitting of A given by collapsing all other topological edges of $A|G_{t_0}$. We must show that for $t > t_0$ the graph $A|G_{t_0}$ has a topological edge that contains a legal segment of length ≥ 2 and that the corresponding splitting coincides with the original.

For this argument it is convenient to work in the unprojectivized Outer space and use the natural parametrization of the folding path, which we rename H_t , so that all folds have speed 1. We also take G_{t_0} as the starting




Figure 6: This is N'_ϵ for the example in Figure 3 assuming that the leftmost and the rightmost widgets are essential.

point H_0 . Thus after time τ the volume of H_τ is $\leq 1 - \tau$ (equality holds if there is only one illegal turn the whole way, otherwise the volume decreases faster). In particular, $\tau < 1$. The legal segment in $A|H_0$ of length $L \geq 2$ may lose a piece of length τ at each end, so we have a legal segment of length $\geq L - 2\tau$ in a topological edge in $A|H_\tau$. After rescaling by $\geq \frac{1}{1-\tau}$ to bring H_τ to volume 1, the legal segment has length $\geq \frac{L-2\tau}{1-\tau} \geq L$ when $L \geq 2$. Moreover, the midpoint of the legal segment is never identified with any other point, so the splitting it determines is independent of τ . $\square$

Corollary 3.2. *Let G_t be a folding path in $\mathcal{X}$. For any f.g. subgroup A of $\mathbb{F}_n$ the image of $A|G_t$ is a reparametrized quasi-geodesic in $\mathcal{S}(A)$ with uniform constants.*

Proof. Progress in $\mathcal{S}(A)$ is made only on the middle interval, and on this interval $A|G_t$ is a folding path in Outer space of A (after rescaling to achieve volume 1, and reparametrizing the path so it is unit speed). The statement then follows from the fact that folding paths project to reparametrized quasi-geodesics. $\square$

Let G_t be a folding path. We say that a collection of subgraphs $H_t \subset G_t$ is *forward invariant* if for every $t < t'$ the folding map $G_t \rightarrow G_{t'}$ takes H_t into $H_{t'}$.

We will denote by Θ_t the image of $A|G_t \rightarrow G_t$, by $\Omega_t \subset G_t$ the set of points with at least two preimages in $A|G_t$, and by $\tilde{\Omega}_t \subset A|G_t$ the preimage of Ω_t in $A|G_t$.

From now on we restrict G_t to the middle segment of Lemma 3.1.

Proposition 3.3. *Assume G_t consists only of the middle segment.*

- Θ_t is a forward invariant core graph whose rank never decreases, and it may increase a bounded number of times,
- both Ω_t and $\tilde{\Omega}_t$ are forward invariant and have a bounded number of components,
- both Ω_t and $\tilde{\Omega}_t$ can change their homotopy type only a bounded number of times, each time by one of the following events (more than one event may occur at the same time):
 - the rank of a component increases, or two components merge into one,
 - an isolated point is created. In this case the rank of Θ_t increases.

Proof. Let $\beta = t_0 < t_1 < \dots < t_M = \gamma$ be the subdivision of $[\beta, \gamma]$ as in Section 2 with the t_i 's corresponding to the “events” in the folding path. First consider the case $t \in (t_i, t_{i+1})$. The image Θ_t is a core graph and locally near a vertex as in Figure 2 it could either be empty or it contains the edge pictured to the left of the vertex as well as at least one edge to the right. The quality of the image does not change as t increases in (t_i, t_{i+1}) .

The subgraph Ω_t near the vertex above could be empty, or it could contain the left edge and 0 or more right edges. Its quality also does not change in the open interval $[t_i, t_{i+1})$. See Figure 7.




Figure 7: Ω_t is locally just the left edge

Thus the homotopy type of Θ_t and of Ω_t is constant for $t \in [t_i, t_{i+1})$ and can change only at the discrete times $t = t_i$, and the same is true for $\tilde{\Omega}_t$. Now let us analyze a small interval $[t_i - \epsilon, t_i]$ as in Section 2. Take a small connected neighborhood N of a vertex in G_{t_i} and look at its preimage N_ϵ in $G_{t_i - \epsilon}$. It is a tree with the additional structure of a height function and widgets. Observe the following:

- If $\Theta_{t_i - \epsilon} \cap N_\epsilon = \emptyset$ then $\Theta_{t_i} \cap N = \emptyset$.
- If $\Theta_{t_i - \epsilon} \cap N_\epsilon \neq \emptyset$, at $t = t_i$ all the components of this intersection merge together. So if the intersection is connected the image does not locally change the homotopy type at $t = t_i$ and otherwise the Euler characteristic decreases, and the change amounts to attaching 1-cells.
- If $\Omega_{t_i - \epsilon} \cap N_\epsilon \neq \emptyset$, at $t = t_i$ all the components of the intersection merge together. So if the intersection is connected Ω_t does not locally change the homotopy type and otherwise the Euler characteristic decreases, and the change amounts to attaching 1-cells.
- If $\Omega_{t_i - \epsilon} \cap N_\epsilon = \emptyset$ but $\Theta_{t_i - \epsilon} \cap N_\epsilon \neq \emptyset$ there are two possibilities. If $\Theta_{t_i - \epsilon} \cap N_\epsilon$ is connected, then this intersection lifts homeomorphically to $A|G_t$ (if two edges of $\Theta_{t_i - \epsilon}$ share a vertex which is not in $\Omega_{t_i - \epsilon}$, their unique lifts must also share a vertex). Thus in this case $\Omega_{t_i} \cap N = \emptyset$ as well. The other possibility is that $\Theta_{t_i - \epsilon} \cap N_\epsilon$ is disconnected. In that case the Euler characteristic of the image decreases at $t = t_i$ as noted above, but note that $\Omega_{t_i} \cap N$ is a single point, i.e. the Euler characteristic of Ω_t increases by 1 at $t = t_i$ (the lifted components cannot merge at $t = t_i$ as that would change the homotopy type of $A|G_t$ at $t = t_i$).

The set of $t = t_i$'s where $\chi(\Theta_t)$ decreases is uniformly bounded, and this set contains all events when Ω_t gains an isolated point. The set of additional times when $\chi(\Omega_t)$ decreases is also bounded, since Ω_t is a subgraph (with respect to the natural CW structure) of G_t and has a uniformly bounded number of components. Thus on the intervals between these times the homotopy types of Θ_t and of Ω_t are constant. We now claim that between these times the homotopy type of $\tilde{\Omega}_t$ is also constant.

To prove the claim, we again analyze an interval $[t_i - \epsilon, t_i]$ as above. We may assume that $\Theta_{t_i - \epsilon} \cap N_\epsilon$ and $\Omega_{t_i - \epsilon} \cap N_\epsilon$ are both connected (otherwise a change in the homotopy type of Θ_t or Ω_t occurs at $t = t_i$) and nonempty. In the analysis we will use the fact that we are in the middle interval, so in particular every vertex of $A|G_t$ has at least two gates. Every component of

the preimage of N (a small metric neighborhood of a vertex v in G_{t_i}) is a metric neighborhood of a point of $\tilde{G}_{t_i}$, the A -cover of G_{t_i} . These components are metric neighborhoods of vertices, and when intersected with $A|G_{t_i}$ become metric neighborhoods of points in $A|G_{t_i}$ (or empty). Some of these intersections are contained in the interior of a topological edge (we will view them as degenerate); the others are small metric neighborhoods $\tilde{N}(\tilde{v}_k)$ of vertices $\tilde{v}_k$ of $A|G_{t_i}$ that map to v . We now have a commutative diagram

$$\begin{array}{ccc} \tilde{N}_\epsilon(\tilde{v}_k) & \longrightarrow & \tilde{N}(\tilde{v}_k) \\ \downarrow & & \downarrow \\ N_\epsilon & \longrightarrow & N \end{array}$$

where $\tilde{N}_\epsilon(\tilde{v}_k)$ is the full preimage of $\tilde{N}(\tilde{v}_k)$ in $A|G_{t_i-\epsilon}$, and it is also a component of the preimage of N_ϵ . Note also that $\tilde{N}_\epsilon(\tilde{v}_k)$ carries the structure of widgets, since $\tilde{v}_k$ has at least two gates, and that the left vertical arrow is an embedding. The union of the images of the components $\tilde{N}_\epsilon(\tilde{v}_k)$'s in N_ϵ (including the degenerate components) is $\Theta_{t_i-\epsilon} \cap N_\epsilon$, and pairwise intersections of the images of components are contained in $\Omega_{t_i-\epsilon} \cap N_\epsilon$. It follows that the image of each component intersects $\Omega_{t_i-\epsilon} \cap N_\epsilon$ in a connected set, i.e. $\tilde{\Omega}_{t_i-\epsilon}$ intersected with each component of the preimage of N_ϵ is nonempty and connected (so the degenerate components are contained in $\tilde{\Omega}_{t_i-\epsilon}$). Thus $\tilde{\Omega}_t$ does not change its homotopy type in the limit.

Finally, we argue that $\tilde{\Omega}_t$ has a uniformly bounded number of components. If not, some topological edge E of $A|G_t$ must intersect $\tilde{\Omega}_t$ in a large number of components. Each point in the frontier of the intersection maps to one of boundedly many vertices in G_t , and each vertex has bounded valence, so we find distinct components in $E - \tilde{\Omega}_t$ with overlapping images, contradicting the definition of $\tilde{\Omega}_t$. $\square$

4 Subfactor projections

4.1 Bounding distances in $\mathcal{S}$

If $G \in \mathcal{X}$, we may collapse all edges of G except for one and obtain a splitting. Choosing different edges gives splittings at distance 1 from each other, so we have an equivariant coarse projection map $\mathcal{X} \rightarrow \mathcal{S}$.

Lemma 4.1. *Let $\phi : G \rightarrow G'$ be a difference of markings so that the preimage of a point has cardinality k . Then $d_{\mathcal{S}}(G, G')$, i.e. the distance in $\mathcal{S}$ between the projections of G, G' , is bounded by a linear function of k . Moreover, if G_t is the path induced by ϕ (see Section 2) then the image of the whole path in $\mathcal{S}$ is also bounded by a linear function in k .*

Proof. G_t is the concatenation of a linear path in a simplex and a folding path. We may assume that the linear path is trivial. By perturbing, we may also assume the point $x' \in G'$ is not a vertex (perturbing slightly can increase cardinality by at most a bounded factor), and that $\phi^{-1}(x')$ does not contain a vertex. If $k = 1$ the two points $x' \in G'$ and $x = \phi^{-1}(x')$ define the same splitting (collapse the complements of small neighborhoods of x, x' on which ϕ is a homeomorphism). The same holds x_t , the preimage of x' in G_t . Otherwise, subdivide G_t so that on the first portion the preimage consists of k points, and on the second portion it consists of $< k$ points. Then the second portion has bounded image by induction, and on the first portion the case $k = 1$ applies. $\square$

Lemma 4.2. *Suppose $\phi : G \rightarrow G'$ is optimal and there are proper core graphs $H \subset G$ and $H' \subset G'$ so that $\phi(H) \subset H'$ and $\phi : H \rightarrow H'$ is a homotopy equivalence. If there is a point $x' \in H'$ such that $\phi^{-1}(x') \subset H$ then $d_{\mathcal{S}}(G, G')$ is bounded.*

Proof. Fold H until $H = H'$ and ϕ is identity on H . This process does not affect edges in the complement of H so there is no movement in $\mathcal{S}$. At the end $\phi^{-1}(x')$ consists of a single point. Now apply Lemma 4.1. $\square$

Remark 4.3. A similar argument shows that if $\phi^{-1}(x') - H$ has k points then the distance in $\mathcal{S}$ between G and G' is bounded by a linear function of k .

4.2 Restricting progress in $\mathcal{S}$ further

Recall that a subgroup A of $\mathbb{F}_n$ is *malnormal* if $gAg^{-1} \cap A \neq \{1\}$ implies $g \in A$. For example, free factors are malnormal. The next lemma shows that if A is malnormal, $A|G_t$ can be making progress only if $\Omega_t \neq G_t$ (see Proposition 3.3).

Lemma 4.4. *Let A be a malnormal f.g. subgroup of $\mathbb{F}_n$. In the setting of Lemma 3.1 assume that for some $t_0 \in [\beta, \gamma)$ (the middle interval) every edge e of G_{t_0} is covered by $A|G_{t_0}$ at least twice, i.e. $\Omega_{t_0} = G_{t_0}$. Then the image of $A|G_t$ for $t \in [t_0, \gamma)$ in $\mathcal{S}(A)$ is a uniformly bounded set, and so is the image of G_t in $\mathcal{S}$ on the same interval.*

Proof. Fix $t \geq t_0$ and let x be the midpoint of an edge of G_t of length

$\geq \frac{1}{3n-3}$. Consider the commutative diagram

$$\begin{array}{ccc} A|G_{t_0} & \rightarrow & A|G_t \\ \downarrow & & \downarrow \\ G_{t_0} & \rightarrow & G_t \end{array}$$

which exists because we are assuming t_0 is not in the first interval $[\alpha, \beta)$, so that $A|G_{t_0}$ maps to the core $A|G_t$. Taking the preimage of x in each of the 4 spaces yields

$$\begin{array}{ccc} R & \rightarrow & P \\ \downarrow & & \downarrow \\ Q & \rightarrow & x \end{array}$$

Because we are assuming that t is in the middle interval $[\beta, \gamma)$ the cardinality of P is uniformly bounded. We will argue:

Claim 1. The cardinality of the preimage R_q in R of any $q \in Q$ is uniformly bounded.

Claim 2. The cardinality of the preimage Q of x in G_{t_0} is uniformly bounded.

The claims together imply that R is uniformly bounded. Lemma 4.1 plus Claim 2 imply that the path from G_{t_0} to G_t makes bounded progress in $\mathcal{S}$ and the boundedness of R implies (by applying Lemma 4.1 to any point in P , which is nonempty by assumption) that the path from $A|G_{t_0}$ to $A|G_t$ makes bounded progress in $\mathcal{S}(A)$.

Proof of Claim 1. From the diagram we have a map $R_q \rightarrow P$. Since P is bounded it suffices to argue that this map is injective. Suppose $a, b \in R_q$ are distinct but map to the same point $p \in P$. Choose an immersed path τ from a to b in $A|G_{t_0}$. Since $(A|G_{t_0}, a) \rightarrow (A|G_t, p)$ is an isomorphism in π_1 , there is a loop τ' in $A|G_{t_0}$ based at a that maps to a loop in $A|G_t$ homotopic to the image of τ , rel p and the same holds in G_t with basepoint x . Since $(G_{t_0}, q) \rightarrow (G_t, x)$ is an isomorphism in π_1 , the images of τ and τ' have to be homotopic rel q . Since they are both immersed, they must be equal, implying that $\tau = \tau'$, contradiction. Alternatively, argue that the folding process does not identify distinct points of R_q .

Proof of Claim 2. Fix an edge e of G_{t_0} . We will argue that e intersects Q in a bounded set. Since the number of edges is bounded, this will prove the claim.

Let u, v be two distinct lifts of e to $A|G_{t_0}$ (they exist by assumption). Define a function $\sigma : e \cap Q \rightarrow P^2$ as follows. If $y \in e \cap Q$, let y_u, y_v be the lifts of y to u, v respectively and let $p(y_u), p(y_v)$ be their images in P .

Define $\sigma(y) = (p(y_u), p(y_v)) \in P^2$. Since P is uniformly bounded, it suffices to argue that σ is injective.

Assume $\sigma(y) = \sigma(y')$ for $y \neq y'$. Let I be the segment of e whose endpoints are y and y' and let I_u, I_v be the lifts of I to u, v respectively. Then the images of I_u and I_v are loops in $A|G_t$ based at distinct points of P (see the proof of Claim 1) and they are lifts of the image of I . But this violates the malnormality of A . $\square$

4.3 Subgroups adjoined in a graph

We say that a f.g. subgroup $A < \mathbb{F}_n$ is *adjoined* in a graph $G \in \mathcal{X}$ if the map $A|G \rightarrow G$ is 1-1 over some edge of G . If we use the notation from Proposition 3.3 and denote by $\Omega \subset G$ the subgraph over which $A|G \rightarrow G$ is at least 2-1 and by $\tilde{\Omega}$ its preimage in $A|G$, then the condition amounts to saying $\tilde{\Omega} \neq A|G$. In other words, G is then obtained from some base graph K (equal to Ω if $A|G \rightarrow G$ is onto) by gluing $A|G$ along a proper subgraph $\tilde{\Omega} \subset A|G$ via an immersion $f : \tilde{\Omega} \rightarrow K$, and the cardinality of $f^{-1}(f(x))$ is ≥ 2 for all $x \in K$.

We say that A is *nearly embedded* in G if $\tilde{\Omega}$ is a forest, and it is *pinched* if $\tilde{\Omega}$ is a finite set. We say A is *embedded* in G if $A|G \rightarrow G$ is 1-1 (i.e. $\tilde{\Omega} = \emptyset$).

Lemma 4.5. *If A is nearly embedded in G then A is a free factor.*

Proof. Let $p : A|G \rightarrow G$ be the canonical immersion. Enlarge the forest $\tilde{\Omega}$ to a maximal tree T for $A|G$ and let E be the set of edges of $A|G$ not in T . Since edges of E aren't in $\tilde{\Omega}$, E is in a 1-1 correspondence with the set of edges $p(E)$ in G . Let $x \in T$ and consider performing Stallings folds to the morphism

$$(T \cup E) \vee_{x=p(x)} (G - p(E)) \rightarrow G$$

Folding the tree T into $G - p(E)$ doesn't affect the homotopy type and when finished the resulting map is 1-1 and onto. In particular, the morphism is a homotopy equivalence. Since $T \cup E$ represents A , the lemma follows. $\square$

Proposition 4.6. *For every folding path G_t there is an interval $[\beta, \tau)$ contained in the middle interval $[\beta, \gamma)$ (see Lemma 3.1) so that:*

- A is adjoined in G_t for all $t \in [\beta, \tau)$, and
- both complementary components of $[\beta, \tau)$ in $[\alpha, \omega]$ have uniformly bounded projections in $\mathcal{S}(A)$.

Thus informally, a folding path can make progress in $\mathcal{S}(A)$ only while A is adjoined in the middle interval.

Proof. First, by Lemma 3.1 the path makes no progress outside the middle interval $[\beta, \gamma)$. Let $[\beta, \tau)$ be the interval inside $[\beta, \gamma)$ consisting of $t \in [\beta, \gamma)$ such that A is adjoined in G_t . This is an initial interval (possibly empty) by the forward invariance of $\tilde{\Omega}_t$, see Proposition 3.3. It remains to argue that on $[\tau, \gamma)$ the path makes bounded progress in $\mathcal{S}(A)$. On this interval we have $\tilde{\Omega}_t = A|G_t$, so if $A|G_t \rightarrow G_t$ is onto the claim follows from Lemma 4.4.

Otherwise, the image is a core graph Θ_t and we may assume it has constant rank by subdividing into a bounded number of subpaths and working with each subpath separately. The volume of Θ_t is ≤ 1 ; after rescaling so that $\text{vol}(\Theta_t) = 1$ and reparametrizing the path so that folds occur at speed 1, we have a folding path with $A|\Theta_t \rightarrow \Theta_t$ surjective, so we apply the same argument. $\square$

Notation 4.7. If A is adjoined in G , we sometimes use the following notation. The immersion $A|G \rightarrow G$ is denoted p . Enlarge a maximal forest for $\tilde{\Omega}$ to a maximal tree T for $A|G$. The set of edges of $A|G - T$ is the disjoint union of a set $E_{\tilde{\Omega}}$ of edges in $\tilde{\Omega}$ and a set E of edges not in $\tilde{\Omega}$. We have that $\langle T \cup E \rangle * \langle T \cup E_{\tilde{\Omega}} \rangle = A$. Further, $T \cup E_{\tilde{\Omega}}$ maps to $G - p(E)$. Let C denote a free factor of $\langle G - p(E) \rangle$ complementary to $\langle T \cup E_{\tilde{\Omega}} \rangle$. There is a morphism $(T \cup E) \vee (T \cup E_{\tilde{\Omega}}) \vee G_C \rightarrow G$ where G_C represents C .

Proposition 4.8. *Assume $n = \text{rk}(\mathbb{F}_n) \geq 3$. If G is a graph such that $B|G$ is embedded and a free factor A is adjoined, then $d_{\mathcal{F}}(A, B) \leq 4$.*

The proof is organized into several cases. One could give a quick argument if 4 is replaced by a larger number.

Proof. Fix notation as above.

Case 1: $E_{\tilde{\Omega}} \neq \emptyset$. There are two cases. If there is an embedded loop β of $B|G \subset G$ contained in $G - p(E)$ then we have

$$A \searrow \langle T \cup E_{\tilde{\Omega}} \rangle \nearrow \langle G - p(E) \rangle \searrow \langle \beta \rangle \nearrow B$$

Otherwise, choose β in $B|G \subset G$. By our assumption, β is not contained in $G - p(E)$. We have

$$A \searrow \langle T \cup E_{\tilde{\Omega}} \rangle \nearrow \langle T \cup E_{\tilde{\Omega}} \cup \beta \rangle = \langle T \cup E_{\tilde{\Omega}} \rangle * \langle \beta \rangle \searrow \langle \beta \rangle \nearrow B$$

where by $\langle T \cup E_{\tilde{\Omega}} \cup \beta \rangle$ we mean the free factor obtained from $\langle T \cup E_{\tilde{\Omega}} \rangle$ by adding an element representing the conjugacy class of β chosen so that

$$\langle T \cup E_{\tilde{\Omega}} \rangle < \langle G - p(E) \rangle < \langle (G - p(E)) \cup \beta \rangle = \langle G - p(E) \rangle * \langle \beta \rangle$$

is a sequence of free factors. Note that $\langle T \cup E_{\tilde{\Omega}} \cup \beta \rangle$ is proper since $rk\langle T \cup E_{\tilde{\Omega}} \rangle < rkA < n$.

For the rest of the proof, assume $E_{\tilde{\Omega}} = \emptyset$, i.e. $\tilde{\Omega}$ is a forest and T is a maximal tree for $A|G$.

Case 2: $rk(G - p(E)) > 1$. Then we may choose an embedded loop β in $G - p(E)$ which is either contained in $B|G$ or $\beta \cup B|G \neq G - p(E)$ and the complement $(G - p(E)) - (\beta \cup B|G)$ contains a non-separating edge. In either case, $d_{\mathcal{F}}(\langle \beta \rangle, B) \leq 2$. Referring back to the proof of Lemma 4.5, $(T - E) \vee (G - p(E)) \rightarrow G$ is a homotopy equivalence and both $T - E$ (representing A) and β are in the complement of a non-separating edge of $G - p(E)$. So we also have $d_{\mathcal{F}}(A, \langle \beta \rangle) \leq 2$.

Case 3: *There is only one edge in E .* Then $3 \leq rk(G) = 1 + rk(G - p(E))$ and we are in Case 2.

Case 4: *There is $e \in E$ and an embedded loop β in $B|G \subset G$ such that β misses $p(e)$.* Then $A \nearrow \langle G - p(e) \rangle \searrow \langle \beta \rangle \nearrow B$.

Case 5: *Every embedded loop in $B|G$ meets every edge of $p(E)$.* Then B has rank 1. For $e \in E$, $(T \cup (E - e)) \vee (G - p(e)) \vee B \rightarrow G$ is a homotopy equivalence (where say the base point corresponds say to an endpoint of e). If $e' \neq e$ is an edge of E , then $A \searrow \langle T \cup e' \rangle \nearrow \langle (T \cup e') \vee B \rangle \searrow \langle B \rangle$. $\square$

Example 4.9. Consider $\mathbb{F}_4 = \langle a, b, c, d \rangle$, $B = \langle a, c, d \rangle$, $A = \langle abaab, cb, abd \rangle$. Then B is embedded and A is adjoined in the rose R . See Figure 8.




Figure 8: Picture of $A|R$. The circle is $\tilde{\Omega}$.

Since A and B have corank 1, the distance between them is even, and in fact it is 4 (A intersects trivially all conjugates of B).

4.4 Coloring of free factors

We will define a notion of a *color* of a free factor, so that the projection $\pi_A(B)$ is well-defined when A and B have the same color (and $A \neq B$). For example, if the color of A is the image of $H_1(A; \mathbb{Z}/2) \rightarrow H_1(\mathbb{F}_n; \mathbb{Z}/2)$ then there can be no *embedding* $A|G_t \rightarrow G_t$ provided G_t contains a subgraph representing B where A and B have the same color but are distinct. This notion is not sufficient for pinching maps. For example, consider $\mathbb{F}_3 = \langle a, b, c \rangle$, $B = \langle a, b \rangle$, and $A = \langle a, cbc^{-1} \rangle$. If G_t is the rose on a, b, c the map $A|G_t \rightarrow G_t$ is a pinching map, since $A|G_t$ is represented by two loops that map to a and b and a connecting edge that maps to c .

We define the color of a free factor A as the function on $H^1(\mathbb{F}_n; \mathbb{Z}/2)$ that to 0 assigns the $\mathbb{Z}/2$ -homology class of A (i.e. the image of $H_1(A; \mathbb{Z}/2) \rightarrow H_1(\mathbb{F}_n; \mathbb{Z}/2)$) and to a nonzero class x , i.e. to an index 2 subgroup $\mathbb{F}_x \subset \mathbb{F}$, it assigns the $\mathbb{Z}/2$ -homology classes of $A \cap \mathbb{F}_x$ and $uAu^{-1} \cap \mathbb{F}_x$ in $\mathbb{F}_x$ for $u \in \mathbb{F}_n - \mathbb{F}_x$. In other words, represent A as a subgraph of a graph G with $\pi_1(G) = \mathbb{F}_n$ and for a nontrivial double cover $\tilde{G} \rightarrow G$ take the homology classes of the (one or two) components of the preimage of the subgraph. E.g. in the above example, A and B have different colors, since unwrapping c yields lifts with different homology classes. More generally, one can show that if A and B are free factors with B embedded in G and A pinched in G , then either $A = B$ or A and B have different colors.

Even this notion is not sufficient for near-embeddings, that is, there are distinct free factors A and B that have the same color and a graph G so that B is embedded and A is nearly embedded in G . For example, take G to be the rank 3 rose with the “identity” marking, i.e. with loops labeled a, b, c , and let $B = \langle c \rangle$, $A = \langle c[a, b]^2 \rangle$. The commutator is squared to insure that homology classes are equal in double covers.

4.5 Projection is well defined

When A, B are two free factors we define the subset $\pi_A(B) \subset \mathcal{S}(A)$ as the set of 1-edge splittings of A obtained as follows: choose any $G \in \mathcal{X}$ where B is embedded and collapse all edges but one in the graph $A|G$. We are interested in restricting to a collection of pairs A, B “in general position” so that $\pi_A(B)$ is uniformly bounded. It is easy to see that for some pairs A, B (e.g. when $A * B = \mathbb{F}_n$) this set is unbounded and the projection is not well defined.

We will argue that the projection of a free factor to the splitting complex of another free factor is well defined assuming either:

- A and B have the same color, or
- A and B are at distance > 4 in $\mathcal{F}$.

First we make the following remark. Fix a graph $\Gamma_0 \in \mathcal{X}$ so that $B|\Gamma_0$ is a rose and $B|\Gamma_0 \rightarrow \Gamma_0$ is an embedding. If $\Gamma \in \mathcal{X}$ is any graph with $B|\Gamma \rightarrow \Gamma$ an embedding, there is a difference of markings map $\phi : \Gamma_0 \rightarrow \Gamma$ that maps $B|\Gamma_0$ to $B|\Gamma$. Let G_t be the path induced by ϕ (see Section 2). The first portion of the path is contained in a simplex, so it induces a uniformly bounded path in $\mathcal{S}(A)$ for any free factor A . The second portion is a folding path in which B is embedded at all times and for $t < t'$ $B|G_t$ maps into $B|G_{t'}$. Thus to prove that $\pi_A(B)$ is uniformly bounded, it suffices to argue that every folding path G_t along which B is embedded makes uniformly bounded progress in $\mathcal{S}(A)$.

Lemma 4.10. *Suppose A and B are distinct free factors that have the same color. Then the projection $\pi_A(B)$ of B to the splitting complex of A is a uniformly bounded set.*

Proof. Let G_t be a folding path along which B is embedded. We will argue that the progress of $A|G_t$ in $\mathcal{S}(A)$ is uniformly bounded. By Proposition 4.6 we may further assume that A is adjoined in every G_t , i.e. $\tilde{\Omega}_t$ is a proper subgraph of $A|G_t$.

Claim. For every t , every point x of $A|G_t - \tilde{\Omega}_t$ maps to $B|G_t \subset G_t$.

Note first that it suffices to check this for points x in the interior of an edge. If x belongs to a nonseparating edge of $A|G_t$, this follows from the assumption that A and B have the same image in $\mathbb{Z}/2$ -homology (some loop in B must cross the image of x). If x belongs to a separating edge and maps to a point $y \in G_t - B|G_t$, consider the double cover of G_t induced by y and how $A|G_t$ lifts to it. We cut G_t open along y and glue two copies crosswise, and do likewise with $A|G_t$ along x . Visibly, $B|G_t$ lifts to two disjoint copies, while $A|G_t$ is cut into two parts, and one part lifts to one copy of $B|G_t$, while the other lifts to the other. This contradicts the assumption that A and B have the same color. The Claim is proved.

Assuming the statement of the lemma is false, fix $t_1 < t_2 < t_3$ so that A_t makes big progress in $\mathcal{S}(A)$ on both $[t_1, t_2]$ and $[t_2, t_3]$ and the homotopy type of $\tilde{\Omega}_t$ is constant for $t \in [t_2, t_3]$ (see Proposition 3.3). We now have 3 cases. Recall that it is possible that $\tilde{\Omega}_t$ is not a core graph.

Case 1. At $t = t_2$ there is a non-core point $x \in \tilde{\Omega}_t$ that maps to $G_t - B_t$.

Consider the preimage of x in $A|G_{t_1}$. This preimage is contained in $\tilde{\Omega}_{t_1}$ by the Claim. We now observe that every edge (or even a legal path) in

$\tilde{\Omega}_{t_1}$ can map over x at most twice. Thus the preimage of x is a uniformly bounded set and we apply Lemma 4.1 to deduce that the distance in $\mathcal{S}(A)$ between $A|G_{t_1}$ and $A|G_{t_2}$ is bounded, contradicting the choice of $t_1 < t_2$.

Case 2. At $t = t_3$ there is a non-core point $x \in \tilde{\Omega}_t$ that maps to $G_t - B|G_t$.

Then argue in the same way and deduce that the distance in $\mathcal{S}(A)$ between $A|G_{t_2}$ and $A|G_{t_3}$ is bounded.

Case 3. Both Case 1 and Case 2 fail, i.e. non-core points of $\tilde{\Omega}_t$ map to $B|G_t$ for $t = t_2$ and $t = t_3$.

In this case, if $\tilde{\Omega}_{t_3}$ maps to $B|G_{t_3}$, then all of $A|G_{t_3}$ maps to $B|G_{t_3}$ so we must have $A \subset B$ and therefore $A = B$ (same color implies the same rank). Therefore, there must be a point x in the core of $\tilde{\Omega}_{t_3}$ that maps outside $B|G_{t_3}$. The preimage of x in $A|G_{t_2}$ is contained in $\tilde{\Omega}_{t_2}$ by the Claim, and in fact it is contained in the core of $\tilde{\Omega}_{t_2}$ by our assumption that Case 1 fails. Thus we are in the situation of Lemma 4.2 and we deduce that $A|G_{t_2}$ and $A|G_{t_3}$ have bounded distance in $\mathcal{S}(A)$, contradicting the choice of t_2 and t_3 . $\square$

Lemma 4.11. *Suppose that the distance in $\mathcal{F}$ between free factors A and B is > 4 . Then the projection of B to $\mathcal{S}(A)$ is well defined up to a bounded set.*

Proof. This follows from Propositions 4.8 and 4.6, since a folding path along which B is embedded cannot have A adjoined anywhere. $\square$

4.6 Verifying conditions of Theorem 1.3

First we check the first bullet in Theorem 1.3. The key is the construction of a “tame path” joining a graph in a folding path that’s making progress in $\mathcal{S}(A)$ to a graph where A is embedded, and so that along that path free factors far from A and the factors of the same color do not make progress.

Lemma 4.12. *Assume that $n = rk(\mathbb{F}_n) \geq 3$ and that a free factor A is adjoined in G . Then there is a path from G to G' so that A is embedded in G' and the path makes only a uniformly bounded progress in $\mathcal{S}(B)$ if B is a free factor such that either $d_{\mathcal{F}}(A, B) > 4$ or A and B have the same color.*

Proof. Use the conventions of Notation 4.7. In particular, we have a morphism $(T \cup E) \vee (T \cup E_{\tilde{\Omega}}) \vee G_C \rightarrow G$. Folding our morphism gives the (inverse of) the desired path G_t . (If some G_t is not a core graph, then we restrict to the core and normalize so that volume is 1.)

We have to check the progress of $B|G_t$ in $\mathcal{S}(B)$. If there is $e \in E$ such that the image in G of $B|G$ crosses $p(e)$ (we say B is *good* for A if this holds)

then visibly this path doesn't make progress in $\mathcal{S}(B)$ since the splitting of B determined by the preimage of a point $y \in \text{int}(p(e))$ is unchanged along the path. Note that B is good for A if A and B have the same color (since e is nonseparating in $A|G$ it suffices to consider image in $\mathbb{Z}/2$ -homology). B is also good for A if $d_{\mathcal{F}}(A, B) > 4$. Indeed, suppose B does not cross $p(e)$ for any $p \in E$. There are two subcases. If $rk(A) > 1$ choose an embedded loop α in $A|G$ missing some $e \in E$. Then $d_{\mathcal{F}}(A, \langle \alpha \rangle) = 1$ and $d_{\mathcal{F}}(\langle \alpha \rangle, B) \leq 3$ since the images of α and $B|G$ are in a proper subgraph of G . If $rk(A) = 1$ choose any rank 1 factor $\langle \beta \rangle$ of B and note that A and $\langle \beta \rangle$ lie in the same rank 2 factor. $\square$

Corollary 4.13. *Assuming free factors A and B have the same color or are at distance > 4 , in the definition of the projection $\pi_A(B) \in \mathcal{S}(A)$ we may take a graph G in which B is adjoined, rather than embedded.*

Proposition 4.14. *Assume $n = rk(\mathbb{F}_n) \geq 3$. Then there is a constant ξ such that the following holds. Let A, B, C be 3 free distinct factors and assume that for each pair either they have the same color or they are at distance > 4 . Then only one of $d_A(B, C), d_B(A, C), d_C(A, B)$ can be larger than ξ .*

Proof. Assume both $d_A(B, C)$ and $d_B(A, C)$ are large. Consider a folding path G_t , $t \in [\alpha, \omega]$, from a graph where B is embedded to a graph where C is embedded. By Proposition 4.6 there is a long interval $[t_1, t_2]$ along which A is adjoined and where the path makes large progress in $\mathcal{S}(A)$. Thus the portion G_t , $t \in [t_2, \omega]$ following this interval is a folding path from a graph in which A is adjoined to a graph where C is embedded, but whose length is much shorter than for the original path. Now use the fact that $d_B(A, C)$ is large plus Corollary 4.13 and run the same argument to conclude that on some long interval $[t_3, t_4]$ in $[t_2, \omega]$ B is adjoined and makes large progress in $\mathcal{S}(B)$. Thus G_t for $t \in [t_3, \omega]$ is a folding path from a graph where B is adjoined to a graph where C is embedded, and by Corollary 3.2 it has a bounded image in $\mathcal{S}(C)$. But this contradicts the fact that on $[t_3, t_4]$ it makes large progress in $\mathcal{S}(C)$. $\square$

We now verify the second bullet of Theorem 1.3.

Proposition 4.15. *Again assume $n \geq 3$. There is a constant ξ' such that the following holds. For any two free factors A, B there are only finitely many free factors C so that*

- A and C have the same color or $d_{\mathcal{F}}(A, C) > 4$,

- B and C have the same color or $d_{\mathcal{F}}(B, C) > 4$, and
- $d_C(A, B) > \xi'$.

We note that the first two bullets are needed so that $d_C(A, B) = \text{diam } \pi_C(A) \cup \pi_C(B)$ is well defined.

Proof. Fix a folding path G_t between a graph where A is embedded and a graph where B is embedded. If $d_C(A, B)$ is large, there is a middle interval of length ≥ 1 (say) along which $C|_{G_t}$ has bounded volume. The number of immersions to a fixed graph Γ with bounded volume is bounded in terms of the injectivity radius of Γ . It follows that only a bounded number of the middle intervals for different C 's can contain the same point, so the number of these intervals is finite. $\square$

5 Consequences

Theorem 5.1. *$Out(\mathbb{F}_n)$ acts on a finite product $Y = Y_1 \times \cdots \times Y_k$ of hyperbolic spaces so that every exponentially growing automorphism has positive translation length.*

In the proof we need the following theorem, announced by Handel-Mosher [7]. Recall [4] that every EG stratum of a $\Phi \in Out(\mathbb{F}_n)$ has an associated *attracting lamination* Λ , carried by a unique free factor $Supp(\Lambda)$ of $\mathbb{F}_n$. The lamination Λ has uncountably many recurrent leaves and none of them are carried by a proper free factor of $Supp(\Lambda)$.

Theorem 5.2 (Handel-Mosher). *Suppose Φ has an EG stratum whose associated lamination Λ has full support, i.e. $Supp(\Lambda) = \mathbb{F}_n$. Then Φ acts with positive translation length on the splitting complex $\mathcal{S}$.*

Proof of Theorem 5.1. Let Y_i be the hyperbolic space produced by Theorem 1.3 for the collection of free factors colored by color i . Then $Out(\mathbb{F}_n)$ acts on the product $Y_1 \times \cdots \times Y_k$.

Let $\Phi \in Out(\mathbb{F}_n)$ be exponentially growing. Choose an EG stratum and let Λ be the associated attracting lamination. There is a smallest free factor A that carries Λ and A is Φ -periodic [4]. Thus for some $j > 0$ we have $\Phi^j(A) = A$, so the restriction of Φ^j to $\mathcal{S}(A)$ has positive translation length, by Theorem 5.2. It follows that Φ^j preserves the copy of $\mathcal{S}(A)$ in the suitable coordinate Y_i and has positive translation length. $\square$

Recall that we have a coarse projection π_A defined on the complement of the 4-ball around $A \in \mathcal{F}$ with values in $\mathcal{S}(A)$. We claim π_A is uniformly coarsely Lipschitz. Indeed, if $B_1 < B_2$ choose a graph $G \in \mathcal{X}$ so that both B_1 and B_2 are embedded in G . We conclude that $\pi_A(B_1) \cap \pi_A(B_2) \neq \emptyset$, which proves our claim.

Theorem 5.3 (Bounded geodesic image theorem). *Let (A_i) be a geodesic in $\mathcal{F}$ and $A \in \mathcal{F}$ at distance > 4 from each A_i . Then the set of projections of the A_i 's to $\mathcal{S}(A)$ is a uniformly bounded set.*

Proof. It suffices to prove the theorem for finite geodesics. We will use the fact that folding paths in $\mathcal{X}$ project to uniform reparametrized quasi-geodesics in $\mathcal{S}$ (see the appendix). There is a folding path G_t whose image in $\mathcal{F}$ is a uniform distance, say K , from (A_i) . Since the projection (where defined) from $\mathcal{F}$ to $\mathcal{S}(A)$ is Lipschitz, we may cut out a portion of (A_i) at distance $\leq K+4$ from A and argue that the projections of the remaining (one or two) geodesics are bounded. Again using that the projection is Lipschitz we may replace the A_i 's by the image of (one or two) folding paths. But a folding path doesn't make progress in $\mathcal{S}(A)$ unless it projects (in $\mathcal{F}$) close to A , by Propositions 4.6 and 4.8. $\square$

6 Final remarks

6.1 Failure of quasi-isometric embedding

We sketch two examples showing that orbit maps $Out(\mathbb{F}_n) \rightarrow Y$ in Theorem 5.1 are not quasi-isometric embeddings.

First, we claim that any polynomially growing automorphism Φ has bounded orbits in Y . By [2] this is equivalent to the assertion that a folding path from Γ to $\Gamma\Phi^m$ makes uniformly bounded progress in every $\mathcal{S}(A)$, where Γ is a relative train track representative for Φ (see [5]). A folding path G_t can be chosen to preserve the stratification. Now consider $A|G_t$ on the middle segment. This segment can be subdivided into a bounded number of subsegments on which the image of $A|G_t \rightarrow G_t$ has constant rank. On each subsegment the splitting of A induced by an edge of $A|G_t$ that maps to the highest stratum of the image is unchanged along the whole subsegment, and this proves the claim.

The second example, pointed out to us by Ursula Hamenstädt, consists of a sequence of automorphisms that are not powers of one automorphism. Take $n = 3$ and $\phi_i : \langle a, b, c \rangle \rightarrow \langle a, b, c \rangle$ given by $a \mapsto a, b \mapsto b, c \mapsto cw_i$, where w_i is an arbitrary (complicated) element of $\langle a, b \rangle$. If the w_i 's get longer, the

images Φ_i of ϕ_i in $Out(\mathbb{F}_3)$ go to infinity, but an argument similar to the above shows that the natural folding path from the “identity rose” R to $R\Phi_i$ does not make progress in any $\mathcal{S}(A)$.

6.2 The case of $\mathcal{F}$

One may replace the splitting complexes $\mathcal{S}(A)$ by the complexes of free factors $\mathcal{F}(A)$ throughout the paper. A folding path G_t can make progress in $\mathcal{F}(A)$ only on the middle interval in Lemma 3.1. In addition, now one can assert that there is only bounded progress when $\tilde{\Omega}_t$ is not a forest. Rephrasing Lemma 4.4, if A is malnormal there is a subinterval of the middle interval where A is nearly embedded and outside this interval progress in $\mathcal{F}(A)$ is uniformly bounded. Using Lemma 4.5 we see that when A is malnormal but not a free factor, the projection $\pi_A^{\mathcal{F}}(B) \subset \mathcal{F}(A)$ is defined for every free factor B and all these sets are contained in a uniformly bounded subset of $\mathcal{F}(A)$, in other words A “comes” with a free factor, canonical up to a bounded distance in $\mathcal{F}(A)$ (assuming $rk(A) > 1$ and in case $rk(A) = 2$ we modify the definition of $\mathcal{F}(A)$ as usual so that it is the Farey graph).

Returning to the case of free factors, Main Theorem holds with projections in $\mathcal{F}(A)$ without any change. We thus get an isometric action of $Out(\mathbb{F}_n)$ on a product of hyperbolic spaces. This time, an automorphism has positive translation length if it has a *bottom* EG stratum.

Likewise, the Bounded geodesic image theorem holds without change after replacing $\mathcal{S}(A)$ with $\mathcal{F}(A)$.

Appendix A Folding paths and hyperbolicity of $\mathcal{S}$

The goal of this appendix is to revisit the Handel-Mosher proof of the hyperbolicity of $\mathcal{S}$ using folding paths. We prove a technical lemma that allows us to use (the usual) folding paths with ≥ 2 gates at every vertex and this simplifies some of the details of the proof. We will work with *liberal folding paths* that include the original Stallings paths [15], the *greedy* folding paths as used in this paper and in [3], and the Handel-Mosher folding paths. The consequence is that liberal folding paths project to reparametrized quasi-geodesics in $\mathcal{S}$. Our proof follows closely the strategy in [7], but it is relatively short partially because we don’t need to be arranging 3 gates condition, and partially because we don’t provide every detail. We also note that Hilion-Horbez [8] give a proof of the hyperbolicity of $\mathcal{S}$ using the strategy of [7], but using Hatcher’s surgery paths in the sphere complex and also find simplifications from this point of view.

A.1 Liberal folding paths

In this appendix, we work with the barycentric subdivision $\mathcal{S}'$ of $\mathcal{S}$. A vertex of $\mathcal{S}'$ can be thought of as a minimal nontrivial simplicial $\mathbb{F}_n$ -tree T with trivial edge stabilizers, or alternatively as the Bass-Serre orbit space $G = T/\mathbb{F}_n$, and we will freely switch between these two points of view. A k -simplex of $\mathcal{S}'$ is a chain $T_0 \rightarrow T_1 \rightarrow \cdots \rightarrow T_k$ of maps between vertices of $\mathcal{S}'$ where each $T_i \rightarrow T_{i+1}$ is obtained by equivariantly collapsing some non-empty set of edges. If $\mathcal{C}$ denotes the space of metric, simplicial $\mathbb{F}_n$ -trees with trivial edge stabilizers (up to equivariant isometry) then there is a surjective forgetful map $\mathcal{C} \rightarrow \mathcal{S}'^{(0)} \subset \mathcal{S}'$. If $T \mapsto S$, we say T represents S .

The *natural CW-structure or triangulation on $G = T/\mathbb{F}_n$* is the one induced by the natural CW-structure on T . We follow [7] and, when working with a specific (probably not natural) triangulation refer, to edges as *edgelets*.

Recall that a *morphism* between two metric trees is an equivariant map which is an isometric embedding on each edgelet for some triangulation of the source. Note that every map between trees that embeds every edge becomes a morphism after choosing a metric in the target and pulling it back. It is always possible to choose a metric so that after suitable subdivisions the map is simplicial and all edges have the same length. Thus our category is morally equivalent to the Stallings category [15].

By an *optimal* morphism we mean a morphism with ≥ 2 gates at every vertex. If $T, T' \in \mathcal{C}$ so that every element of $\mathbb{F}_n$ which is elliptic in T is also elliptic in T' , then there is an optimal morphism $T'' \rightarrow T'$ where T'' is obtained from T' by equivariantly collapsing some edges and possibly changing the metric.

We allow *liberal folding*, i.e. illegal turns can be folded with any speed, folding can pause and then continue, and we even allow not doing anything on a subinterval. More precisely, a *(liberal) folding path guided by optimal f* is

- a path G_t , $t \in [0, L]$, in $\mathcal{C}$ and a collection $\{G_{t_1} \rightarrow G_{t_2}\}_{0 \leq t_1 < t_2 \leq L}$ of morphisms


such that

- $G_0 \rightarrow G_L$ is f and $G_{t_1} \rightarrow G_{t_2} \rightarrow G_{t_3}$ is a factorization of $G_{t_1} \rightarrow G_{t_3}$ for all $0 \leq t_1 < t_2 < t_3 \leq L$.

We occasionally use $[G_{t_1}, G_{t_2}]$ to denote $\{G_t \mid t \in [t_1, t_2]\}$. The proof of [3, Proposition 2.2] shows *mutatis mutandis* that, given f , there is a (greedy) folding path guided by f .

A.2 Collapses and expansions

We will denote trees by $\tilde{G}$ (with various sub and superscripts) and the quotient graphs with G (with the same sub and superscripts). If G_t , $t \in [0, L]$, is a liberal folding path and if $F_L \subset \tilde{G}_L$ is a proper equivariant forest, define $\tilde{G}'_t = \tilde{G}_t / F_t$ where F_t is the preimage of F_L in $\tilde{G}_t$.




Lemma A.1. G'_t is a liberal folding path.

Note that even if G_t is a greedy folding path, the quotient G'_t may not be greedy since we could collapse the parts where folding takes place (so in fact G'_t may have subintervals on which it is constant).

Proof. There is a natural map $f' : \tilde{G}'_0 \rightarrow \tilde{G}'_L$ with respect to which $\tilde{G}'_t$ is a liberal folding path guided by f' , but we must show that every vertex has at least 2 gates. See [7, Proposition 4.4]. This is clear except possibly at vertices coming from the components of F_0 . If d, d' are distinct directions coming out of one such component C and if $d' = g(d)$ for some $g \in \text{Stab}(C)$, then d, d' must map to distinct directions out of $f(C)$ since direction stabilizers are trivial. Thus we are reduced to considering the case when $\text{Stab}(C) = 1$, i.e. when C is a nondegenerate finite tree. If all directions out of C map to the same direction out of $f(C)$, say based at $v \in f(C)$, then the finite nondegenerate tree $f(C)$ has an endpoint w distinct from v . Any vertex of C that maps to w can have only one gate, contradiction. $\square$

There is a reverse operation of *expansion*. Let G_t , $t \in [0, L]$, be a folding path and let $\tilde{G}'_L \rightarrow \tilde{G}_L$ be a collapse map. Define $\tilde{G}'_t$ as the pull-back of the maps $\tilde{G}_t \rightarrow \tilde{G}_L$ and $\tilde{G}'_L \rightarrow \tilde{G}_L$. The induced map $\tilde{G}'_t \rightarrow \tilde{G}'_L$ is an isometry on subtrees that map to point in $\tilde{G}_t$.



As t increases the path G'_t is a liberal folding path except at times t when in the path G_t distinct preimages of points in G_L that got blown up

get identified under folding. At those times different copies of blow-up trees get identified. To transform G'_t into a folding path insert an interval where these different copies get identified (equivariantly) by folding. Thus the blow-up operation is not completely canonical as it requires a choice (the choice could be made canonical by taking the greedy identification path).

Lemma A.2. *Thus modified, G'_t is a liberal folding path.* $\square$

Thus given a folding path and an expansion of the terminal graph, there is an induced folding path. We will normally suppress the required insertions of constant paths and reparametrizations.

A.3 Distance estimates in $\mathcal{S}'$

With little change, the proof of Lemma 4.1 shows:

Lemma A.3. *Let $f : G \rightarrow G'$ be an optimal morphism. Then $d_{\mathcal{S}'}(G, G')$ is bounded by a linear function of the cardinality of $f^{-1}(y)$ for any $y \in G'$.*

In particular, if G_t is a folding path guided by f then the diameter of the image of the folding path in $\mathcal{S}'$ is bounded by a linear function of the cardinality of $f^{-1}(y)$ for any $y \in G'$.

In the next lemma we will consider finite sets R and B of red and blue points in G' . We will assume $R \cap B = \emptyset$ and also that $R \cup B$ is disjoint from the vertex set. A *mixed region* in G' is a complementary component of $R \cup B$ whose frontier intersects both R and B . By pulling back red and blue points, we obtain red and blue points in G , and we can talk about mixed regions in G . The following lemma is a restatement of [8, Proposition 6.2] in the setting of optimal morphisms.

Lemma A.4. *Let $f : G \rightarrow G'$ be an optimal morphism. Suppose R and B are nonempty disjoint finite sets in G' also disjoint from the set of natural vertices. Then $d_{\mathcal{S}'}(G, G')$ is bounded by a linear function of the number of mixed regions in G .*

This generalizes Lemma A.3, e.g. take $R = \{y\}$ and B is a single nearby point. In practice, R and B will be chosen so that $R \cup B$ intersects every edge in one point, and then the complementary regions are trees, and in this form the lemma appears in [7]. Note that bounding the number of mixed regions in G amounts to saying that an edge of G can map over both colors many times, but the number of times it goes from one color to the other is bounded.

Proof. We subdivide G' so that the restriction of f to each edge (i.e. edgelet) of the induced subdivision of G is an isometry and each edgelet of the subdivision of G' contains at most one colored point. Thus each edgelet is colored blue, red, or neutral. By collapsing all neutral edgelets, we may assume there are only red and blue edgelets. In particular, mixed regions correspond to *mixed vertices*, i.e. vertices adjacent to both red and blue edgelets. Consider a folding path G_t , $t \in [0, L]$, guided by f . We impose the requirement that the folding path is obtained by iteratively folding only red edgelets for as long as possible; and then only folding blue edgelets for as long as possible, etc. Thus the folding path is broken up into segments, called *phases*, and we talk about red and blue phases, depending on which color is allowed to get identified. The proof now follows from three observations:

- Under folding the number of mixed vertices never goes up; in fact the complexity $(N, -r)$ can never increase, where N is the number of mixed vertices and r is the sum of the ranks of the stabilizers over the orbits of mixed vertices (viewing G_t now as trees). More precisely, the complexity changes only in the event that two mixed vertices merge into one – if they are in different orbits N decreases, and if they are in the same orbit r increases.
- The progress in $\mathcal{S}'$ is bounded during any phase. E.g. during a red phase, blue edgelets define splittings that don't change during this phase.
- Complexity has to decrease at least once during each phase, except possibly first and last.

To prove the third bullet, consider a phase, say red, starting with G_t . Viewing G_t as a tree, it is decomposed into maximal red and blue subtrees. If our red phase is not the first phase, each component of the blue subforest of G_t embeds in G_L . If our red phase is not the last, then there is a first time during the phase when a red fold identifies points in distinct blue subtrees and hence complexity decreases. $\square$

Definition A.5. A *hanging tree* is a tree with a base vertex v and a train track structure in which v and all valence one vertices have only one gate, and every other vertex has two gates, with the direction towards v being its own gate. We draw a hanging tree with v on top and call it the top vertex.

Example A.6. Consider the map $f : R \rightarrow R$ where R is the rank 3 rose with the identity marking given by $x \mapsto x$, $y \mapsto x^k y x^{-k}$ and $z \mapsto x^k z x^{-k}$. Then

f can be thought of as obtained from the identity $R \rightarrow R$ by “rotating” k times around x . The preimage of a point in x is contained in the loop x together with the hanging tree consisting of initial and terminal segments in edges labeled y and z . This is an example where (i) fails in the next lemma.

Lemma A.7. *Let $f : G \rightarrow G'$ be an optimal morphism between graphs in Outer space and suppose that $e \subset G$ and $e' \subset G'$ are edges that define the same splitting. Then one of the following holds:*

- (i) *The cardinality of the preimage of a point $y' \in \text{int } e'$ is uniformly bounded in G .*
- (ii) *There is a hanging tree $H \subset G$ with one endpoint of e attached to the top vertex and the other endpoint to a point of H so that e together with a (possibly degenerate) segment of H forms an embedded legal loop E . The preimage of some $y' \in e'$ is contained in $H \cup e$ and intersects E in one point. Both endpoints of e have two gates.*

Proof. Construct a folding path G_t guided by f so that on an initial segment from G to say G_s the image of e is not involved in any folds and so that the map $G_s \rightarrow G'$ does not admit any folds not involving the image e_s of e . Let $\hat{e}_s$ be the natural edge in G_s that contains the edge e_s . By assumption, there are no illegal turns in G_s except for those involving e_s . Now consider cases.

Case 1. $G_s - \hat{e}_s$ has no valence 1 vertices. Then $G_s - \hat{e}_s \rightarrow G'$ is an immersion and on fundamental groups induces an isomorphism $\pi_1(G_s - \hat{e}_s) \rightarrow \pi_1(G' - e')$ (if $\hat{e}_s$ and e' are separating, we mean an isomorphism for each component). It follows that $G_s - \hat{e}_s \rightarrow G' - e'$ is an embedding (in fact, a homeomorphism, unless e' is a loop connected to the rest of the graph by a separating edge).

Now since $G_s \rightarrow G'$ is a homotopy equivalence, the image of $\hat{e}_s$ crosses e' exactly once. Thus for any point y' in e' the preimage in G_s is a single point y_s in $\hat{e}_s$. Now note that if σ is any edge (or a legal path) in $G - e$, its image in G_s is disjoint from e_s , so it can cross y_s at most twice, and conclusion (i) follows.

Whenever the endpoints of $\hat{e}_s$ are distinct, we are in Case 1. Moreover, we may assume that $\hat{e}_s$ is a loop with vertex a that has valence 3.

Case 2. $\hat{e}_s$ is a loop with vertex a with valence 3, and the two directions into $\hat{e}_s$ are equivalent. Then the turn between the two $\hat{e}_s$ -directions is the only illegal turn in G_s , so G' is obtained from G_s by folding the “monogon” $\hat{e}_s = e_s$. Now take $y' \in G'$ to be a point in the image of e_s . The preimage

of y' has at most two points in G_s and is contained in e_s . The image in G_s of an edge $u \neq e$ in G cannot cross e so (i) again holds.

Case 3. All turns in G_s are legal. Then $G_s = G'$ so (i) holds with y' a point in e_s .

Case 4. $\hat{e}_s$ is a legal loop with vertex a with valence 3 and two gates. Thus there is a separating edge z incident to a . Then the complement of $\hat{e}_s \cup z$ embeds into $G' - e'$ as in Case 1. Now fix $y' \in e'$, and note that the preimage of y' in G_s is contained in $\hat{e}_s \cup z$ and meets $\hat{e}_s$ in a single point. We claim that the preimage of $\hat{e}_s \cup z$ in G is the union of e and a hanging tree, with one endpoint of e attached to the top of the tree, and the other endpoint somewhere on the tree, and so that e is contained in a legal embedded loop. Once established, the claim proves that (ii) holds.

One end of $e_s \subset G_s$ must be involved in the illegal turn. Without loss, say e_s is oriented so that the terminal direction is involved in the illegal turn. If u is any point of $z \cup \hat{e}_s - e$ there is a unique embedded legal path P_u starting at u and ending at the initial vertex of e_s . If $x \in G$ is any point that maps to some $u \in z \cup \hat{e}_s - e$ (except the endpoint of z not in e_s), using the fact that $G \rightarrow G'$ is optimal we see that there is a unique legal path Q_x in G that maps isometrically to P_u and ends at the initial vertex of e_s (one can lift P_u using that there are ≥ 2 gates everywhere; the lift must end at the initial vertex of e since $G \rightarrow G_s$ is 1-1 over e_s ; the lift is unique since otherwise a loop gets killed). The union of all Q_x 's is a hanging tree with the top vertex at the initial vertex of e . $\square$

Remark A.8. If $f : G \rightarrow G'$ is an optimal map and $f(x_1) = f(x_2)$ then x_1 and x_2 are joined by a unique immersed path whose image is nullhomotopic rel endpoints. The existence follows from π_1 -surjectivity of f , and uniqueness from π_1 -injectivity. We call this path a *vanishing path*. Now suppose that we are in the situation of Lemma A.7 and that (ii) holds. If $x_1, x_2 \in f^{-1}(y')$ then the vanishing path between x_1 and x_2 is contained in $H \cup e$. This can be seen by considering the factorization $G \rightarrow G_s \rightarrow G'$ from the proof of Lemma A.7. To elaborate, vanishing paths for points in the preimage of y' in $\hat{e}_s \cup z \subset G_s$ are contained in $\hat{e}_s \cup z$ since the loop $\hat{e}_s$ folds with (a part of) z on its way to G_L and identifies all points in the preimage of y' . Since the map $H \cup e \rightarrow \hat{e}_s \cup z$ is a π_1 -isomorphism, the same argument as above shows that there is a path α in $H \cup E$ connecting x_1 and x_2 whose image (tightened) is the vanishing path between the images of x_i , hence α is vanishing as well.

Note that every immersed path in $H \cup e$ has at most one illegal turn, hence a vanishing path has exactly one illegal turn. If $x_1, x_2 \in f^{-1}(y') - E$ are

connected by a legal segment J , say oriented towards E , then the vanishing path from x_1 to x_2 necessarily has the form $Ju \cdot E^{-k}u^{-1}$ where $\cdot$ denotes the illegal turn. See Figure 9.




Figure 9: Vanishing path in $\hat{e}_s \cup z \subset G_s$

In particular, given any folding path G_t guided by f (not necessarily passing through G_s above), if x_1 and x_2 are identified in some G_t , then they are both identified with $y \in f^{-1}(y') \cap E$ (the vanishing path must completely fold). Consequently, if $x_1, \dots, x_k \in f^{-1}(y')$ are identified in G_t , then either k is bounded, or all these points are identified with y (the hanging tree H has bounded complexity and given a sufficiently large subset, two of the points are connected by a legal path).

Proposition A.9. *Suppose $f : G \rightarrow G'$ is an optimal morphism such that G and G' have edges that determine the same splitting. Then the image in S' of a liberal folding path G_t , $t \in [0, L]$, guided by f is bounded.*

Proof. By blowing up G_L if necessary and applying Lemma A.2, we may assume that G_L is in Outer space, i.e. point stabilizers are trivial. Apply Lemma A.7 to $G_0 \rightarrow G_L$. If (i) holds, then the conclusion follows from Lemma A.3, so suppose (ii) holds. Let t_1 be the first time that the preimage of y' in G_{t_1} consists of a single point. Then for any small $\epsilon > 0$ the paths G_t for $t \in [0, t_1 - \epsilon]$, $t \in [t_1 - \epsilon, t_1]$ and $t \in [t_1, L]$ map to bounded sets in S' . Indeed, this is true for the last one by Lemma A.3, for the middle path by the smallness of ϵ , and for the first path again by Lemma A.3 applied to

any preimage of y' in $G_{t_1-\epsilon}$ not in the image of E (by the last sentence in Remark A.8 the preimage in G_0 of such a point is bounded). $\square$


A.4 Combing diagrams and rectangles

A *combing rectangle* is a pair of folding paths G_t and G'_t together with collapse maps $G_t \rightarrow G'_t$ that commute with folding maps so that the top row is obtained from the bottom row either by collapsing as in Lemma A.1 or by expanding as in Lemma A.2. We represent a folding rectangle by a rectangle with sides oriented to indicate the direction of folding and of collapse. The folding direction is horizontal and the collapsing is vertical. A combing diagram consists of combing rectangles and folding paths. All horizontal rows are (liberal) folding paths and in particular all horizontal maps are optimal.

A *zig-zag path* is an alternating sequence of collapses and expansions. Every geodesic path in $\mathcal{S}'$ is a zig-zag path. We don't insist here on minimal triangulations where all vertices have valence > 2 .


A.5 Simplifying

Here we introduce simplifying moves on combing diagrams. Suppose we have a combing diagram




Let F, F' be the forests in $\tilde{G}_L''$ collapsed by the two maps. Then

- If $F'' = F \cap F'$ contains a nondegenerate segment, there is a combing diagram



where $\tilde{G}_t'''$ is obtained from $\tilde{G}_t''$ by collapsing the components of the preimages of the forest F'' .


- If $F \cup F' \neq \tilde{G}_L''$, there is a combing diagram




where $\tilde{G}_t'''$ is obtained from $\tilde{G}_t''$ by collapsing the preimage of $F \cup F'$.

The following is our version of Handel-Mosher's Proposition 6.5 from [7].

Theorem A.10 (Handel-Mosher). *For every combing diagram*



there is $0 \leq K \leq L$ and a combing diagram



so that the folding paths $[\hat{G}'_K, G'_L]$ and $[G''_K, G''_L]$ are uniformly bounded in S' .

Proof. Equivariantly subdivide $\tilde{G}_L$ into finitely many orbits of edgelets so that each edgelet either collapses or maps isometrically by either vertical

map. Using the simplifying operations from section A.5 and replacing the middle folding path by a quotient, we may assume that every edgelet in G_L collapses under exactly one of the two vertical maps. We will color edgelets in red and blue so that red edgelets collapse under $G_L \rightarrow G'_L$ and blue edgelets under $G_L \rightarrow G''_L$. By further subdividing, we will also assume that

- the preimage of the interior of an edgelet in $\tilde{G}_0$ contains no natural vertices,
- the restriction of $\tilde{G}_0 \rightarrow \tilde{G}_L$ to each component of the preimage of an edgelet is an isometry, and
- blue edgelets of $\tilde{G}_L$ map isometrically to $\tilde{\Gamma}$ via $\tilde{G}_L \rightarrow \tilde{G}'_L \rightarrow \tilde{\Gamma}$.

For $t \in [0, L]$, color G_t by pulling back via $G_t \rightarrow G_L$. In particular, G_t is colored red and blue, $G_t \rightarrow G''_t$ is obtained by collapsing components of blue subforests (so G''_t is colored red), and $G_t \rightarrow G'_t$ is obtained by collapsing red subforests (so G'_t is colored blue). To focus on the colors, we write, for example, G_t^R for G_t with all components of the blue subforests collapsed. So, $G_t^R = G''_t$ and is colored red. Similarly, $G_t^B = G'_t$ is colored blue.


By Lemma A.4 applied to blue and red, there is $K \in [0, L]$ so that $[G_K, G_L]$ is bounded in $\mathcal{S}'$ and a natural edge e of G_K that contains a red arc with at least one blue arc on either side. By further subdividing G_L , we may also assume:

- the preimage in G_K of the interior of an edgelet of G_L contains no vertices.

In particular, G_K is a union of colored edgelets with pairwise disjoint interiors each mapping isometrically to an edgelet of G_L . Let r be a red edgelet in e with at least one blue edgelet on each side in e and so that there is a blue edgelet b in e immediately adjacent to r .

Denote by G_K^{rB} the result of collapsing in G_K all red edgelets except r and by G_K^r the result of additionally collapsing all blue edgelets. For $t \in [0, K)$, G_t^{rB} and G_t^r are obtained as above by collapsing components of

preimages of edgelets. We have the combing diagram:




By assumption, there is an equivariant homeomorphism $h : G_K^{rB} \rightarrow G_K^B$ preserving edgelets not in the orbit of $r \cup b$ and mapping $r \cup b$ to b . Roughly, our desired diagram is obtained by deleting the top left element G_0^B from the diagram above. To make this precise, we have to adjust the metric on G_t^{rB} and replace $G_K^{rB} \rightarrow G_K^B$ with an isometry. We now do this.

Let $\hat{G}_K^{rB}$ denote G_K^{rB} with the metric induced by h and let $\hat{G}_0^{rB}$ denote G_0^{rB} with the pullback metric via $G_0^{rB} \rightarrow G_K^{rB} \xrightarrow{h} G_K^B$. Edgelets and colors of $\hat{G}_0^{rB}$ and $\hat{G}_K^{rB}$ are induced by the natural homeomorphisms with respectively G_0^{rB} and G_K^{rB} . We identify r with its image in $\hat{G}_K^{rB}$. By construction, we have morphisms $\hat{G}_0^{rB} \rightarrow \hat{G}_K^{rB} = G_K^B \rightarrow \Gamma$. It remains to check that the composition is optimal.


By construction, our composition is injective when restricted to edgelets. Since $G_0^B \rightarrow \Gamma$ is optimal, a vertex of $\hat{G}_0^{rB}$ with no incident edgelets in the orbit of the preimage of r has two gates. Since r is interior to a natural edge of G_K^{rB} , turns at endpoints of edgelets in r are legal with respect to $\hat{G}_K^{rB} \rightarrow \Gamma$. Hence, we only need to show that an endpoint of an edgelet of $\hat{G}_0^{rB}$ in the orbit of the preimage of b has two gates with respect to $\hat{G}_0^{rB} \rightarrow \hat{G}_K^{rB}$. But this is clear since $G_0^{rB} \rightarrow G_K^{rB}$ is optimal. $\square$

Corollary A.11. *There is a constant $M = M(\text{rank } \mathbb{F}_n)$ so that the following holds. Suppose we are given a combing diagram which is a vertical stack of $N > 1$ combing rectangles with the projection to $\mathcal{S}'$ of the bottom row of diameter $> MN$.*



Then there is a combing diagram consisting of a stack of two rectangles with top and bottom paths extended to folding paths so that:

- the bottom folding path equals the bottom folding path of the given diagram,
- the top right corner equals the top right corner of the given diagram,
- if the original top folding path is extended to the right, the same extension forms a folding path in the resulting diagram,
- the vertical arrows in the two resulting rectangles are down-up,
- paths not included in the rectangles have diameter $< MN$ in $\mathcal{S}'$.



Proof. When $N = 2$ this is Theorem A.10. Proceed by induction, combining consecutive collapse maps into a single collapse map. $\square$

A.6 Verification of Masur-Minsky axioms

To prove that $\mathcal{S}'$ is hyperbolic and that folding paths are reparametrized geodesics it suffices to verify:

- (1) Folding paths form a coarsely transitive family of paths.
- (2) For each folding path G_t there is a coarse Lipschitz retraction $\mathcal{S}' \rightarrow G_t$.
- (3) Every folding path is uniformly strongly contracting.

See [12] and also [7, Section 3].


We now proceed to verify these properties.

A.6.1 Folding paths are coarsely transitive

If $G_1, G_2 \in \mathcal{S}'$ let $G'_i \in \mathcal{S}'$ be a rose obtained from G_i by expanding, then collapsing. After adjusting the metric on G'_1 , there is a folding path from G'_1 to G'_2 .

A.6.2 Definition of the projection.


Let G_t be a folding path and $\Gamma \in \mathcal{S}'$. Following [7], define the *projection* $\pi(\Gamma)$ of Γ to the folding path G_t as the time T (or the graph G_T), where T is the sup of the times t such that there is a combing diagram as pictured where Γ denotes any representative of Γ .



If no such combing diagram exists, the sup is defined to be 0. (It can be shown that it always exists, i.e. there is a collapse followed by an expansion $G_0 \rightarrow R \leftarrow G'_0$ and an optimal map $G'_0 \rightarrow \Gamma$.)

A.6.3 The projection is a coarse retraction


Let $\Gamma = G_{t_0}$ (as underlying graphs, but with possibly different metrics). Then clearly $T = \pi(\Gamma) \geq t_0$, since we can construct combing rectangles from $G_{t_0} \rightarrow R \leftarrow \Gamma$, where R has one edge. Suppose we have a combing diagram as pictured.




We must show that $d_{\mathcal{S}'}(G_{t_0}, G_t)$ is uniformly bounded. Apply Lemma A.9 to the optimal map $G'_{t_0} \rightarrow \Gamma = G_{t_0}$ (since G_{t_0} and G'_{t_0} collapse to the same graph, they each have an edge that determines the same splitting). The conclusion is that the folding path from G'_{t_0} to G'_t has bounded image in $\mathcal{S}'$, so the same is true for the path $[G_{t_0}, G_t]$.

A.6.4 The projection is coarsely Lipschitz

Suppose $\Gamma \rightarrow \Gamma'$ is a collapse. We first observe that $\pi(\Gamma) \geq \pi(\Gamma')$. The argument is in the diagram below. Construct a combing rectangle using the folding path represented by the horizontal line ending in Γ' and the collapse map $\Gamma \rightarrow \Gamma'$ (note that this may require inserting constant paths and reparametrizing) and combine it with the given combing rectangles. The top two rectangles on the left form one combing rectangle.




Now we must show that the segment $[\pi(\Gamma'), \pi(\Gamma)]$ has a uniformly bounded projection in $\mathcal{S}'$. This follows from Corollary A.11 with $N = 3$ applied to the stack of 3 rectangles in the diagram below.




A.6.5 Folding paths are uniformly strongly contracting

Let G_t be a folding path and $\Gamma \in \mathcal{S}'$ so that $d_{\mathcal{S}'}(\Gamma, G_t) > R$ for every t and R is large. Suppose $\Gamma' \in \mathcal{S}'$ is such that $d_{\mathcal{S}'}(\Gamma, \Gamma') < \mu R$ for a certain $\mu > 0$. After possibly adjusting the metric on Γ' there is a zig-zag path from Γ to Γ' of length $< \mu R$. Take a combing diagram associated with the projection of Γ to G_t and add to it the stack of combing rectangles induced by the zig-zag path (in the diagram $d_{\mathcal{S}'}(\Gamma, \Gamma') = 3$).



Apply Corollary A.11 (with N equal to the height of the stack) to replace the stack of rectangles with a stack of height two, plus the trailing folding paths. The assumption on the distances guarantees that this stack continues beyond the combing rectangles for the projection.



Apply Corollary A.11 one more time with $N = 4$ and deduce that the projection of Γ' cannot precede the projection of Γ by more than a bounded amount. Reversing the roles of Γ and Γ' proves the statement.

To summarize, we have

Theorem A.12. $\mathcal{S}'$ (and hence also $\mathcal{S}$) is hyperbolic and liberal folding paths are uniform reparametrized quasi-geodesics.

References

- [1] Jason A. Behrstock. Asymptotic geometry of the mapping class group and Teichmüller space. *Geom. Topol.*, 10:1523–1578, 2006.
- [2] Mladen Bestvina, Kenneth Bromberg, and Koji Fujiwara. Constructing group actions on quasi-trees and applications to mapping class groups. arXiv:1006.1939.
- [3] Mladen Bestvina and Mark Feighn. Hyperbolicity of the complex of free factors. arXiv:1107.3308.
- [4] Mladen Bestvina, Mark Feighn, and Michael Handel. The Tits alternative for $\text{out}(F_n)$. I. Dynamics of exponentially-growing automorphisms. *Ann. of Math. (2)*, 151(2):517–623, 2000.
- [5] Mladen Bestvina and Michael Handel. Train tracks and automorphisms of free groups. *Ann. of Math. (2)*, 135(1):1–51, 1992.
- [6] Stefano Francaviglia and Armando Martino. Metric properties of Outer space. *Pub. Mat.*, 55:433–473, 2011.
- [7] Michael Handel and Lee Mosher. The free splitting complex of a free group I: Hyperbolicity. arXiv:1111.1994.
- [8] Arnaud Hilion and Camille Horbez. The hyperbolicity of the sphere complex via surgery paths. arXiv:1210.6183.
- [9] Ilya Kapovich and Kasra Rafi. On hyperbolicity of free splitting and free factor complexes. arXiv:1206.3626.
- [10] Johanna Mangahas. Uniform exponential growth of subgroups of the mapping class group. *Geom. Funct. Anal.*, 19(5):1468–1480, 2010.
- [11] Brian Mann. Hyperbolicity of the cyclic splitting complex. arXiv:1212.2986.
- [12] Howard A. Masur and Yair N. Minsky. Geometry of the complex of curves. I. Hyperbolicity. *Invent. Math.*, 138(1):103–149, 1999.
- [13] Howard A. Masur and Yair N. Minsky. Geometry of the complex of curves. II. Hierarchical structure. *Geom. Funct. Anal.*, 10(4):902–974, 2000.

- [14] Lucas Sabalka and Dmytro Savchuk. Submanifold projection. arXiv:1211.3111.
- [15] John R. Stallings. Topology of finite graphs. *Invent. Math.*, 71(3):551–565, 1983.